

2020-01-22
2019-2458

FOISON Glufosinate Ammonium 200 SN

GROUP	10	HERBICIDE
-------	----	-----------

FOR SALE FOR USE ON GLUFOSINATE AMMONIUM TOLERANT FIELD CORN AND ON GLUFOSINATE AMMONIUM TOLERANT SOYBEANS IN CANADA AND ON GLUFOSINATE AMMONIUM TOLERANT CANOLA IN EASTERN CANADA AND BRITISH COLUMBIA (INCLUDING THE PEACE RIVER REGIONS)

**SOLUTION
COMMERCIAL**



CAUTION - EYE IRRITANT

REGISTRATION NUMBER: 33386 PEST CONTROL PRODUCTS ACT

ACTIVE INGREDIENT:
glufosinate-ammonium 200 g/L
NET CONTENTS: 1L - 1,000 L

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

Product Information: +1-519-702-2185

**Jiangsu Good Harvest-Weien Agrochemical Co., LTD
Laogang, Qidong City, Jiangsu 226221 China
Canadian representative: Hanson Industry Company
569 Rosehill Ave, Stittsville, ON K2S 0E8
Tel: +1-519-702-2185**

In case of spills, poisoning or fire, telephone emergency response number: +1-519-702-2185

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. CAUTION: EYE IRRITANT. May irritate eyes. Avoid contact with eyes, skin or clothing. Avoid breathing spray mist. Wash thoroughly after using and before eating, drinking or smoking.

Protective Clothing and Equipment: Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. In addition, wear protective eyewear (goggles or face shield) during mixing and loading. Wear a respirator with a NIOSH-approved organic-vapour-removing cartridge with a prefilter approved for pesticides, or a NIOSH-approved canister approved for pesticides during mixing, loading and application.

Re-entry Restrictions: DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. If required, certified applicators may enter treated areas within 24 hours for short-term tasks not involving hand labour if at least 4 hours has passed since application and long-sleeved shirt, long pants, chemical-resistant coveralls, chemical-resistant footwear, socks, goggles, chemical-resistant gloves, hat and a respirator with a NIOSH-approved organic-vapour-removing cartridge with a prefilter approved for pesticides OR a NIOSH-approved canister approved for pesticides is worn. Time spent in the treated area cannot exceed 1 hour in a 12-hour period during the restricted re-entry interval.

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

DO NOT apply using aerial application equipment

If this pest control product is to be used on a commodity that may be exported and you require information regarding Maximum Residue Limits for an importing country, please contact Jiangsu Good Harvest-Weien Agrochemical Co., Ltd. at +1-519-702-2185.

FIRST AID: Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention. If swallowed, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person. **If in eyes,** hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. **If on skin or clothing,** take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice. **If inhaled,** move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

TOXICOLOGICAL INFORMATION:

Note to Physician: If this product is ingested, endotracheal intubation and gastric lavage should be performed as soon as possible, followed by charcoal and sodium sulfate administration.

ENVIRONMENTAL PRECAUTIONS: TOXIC to aquatic organisms and non-target terrestrial plants. Observe buffer zones specified under Directions for Use. To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body. Avoid spray drift onto susceptible plants and USE ONLY FOR RECOMMENDED PURPOSES AND AT RECOMMENDED RATES. DO NOT APPLY BY AIR.

STORAGE: CANNOT be stored below freezing. If stored for 1 year or longer, shake well before using. Store the tightly closed container away from feeds, seeds, fertilizers, plants and foodstuffs. Do not use or store in or around the home. Keep in original container during storage.

DISPOSAL: Recyclable Container Disposal: Do not reuse this container for any purpose. This is a recyclable container and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site: 1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank. 2. Make the empty, rinsed container unsuitable for further use. If there is no container collection site in your area, dispose of the container in accordance with provincial requirements. **Returnable Container Disposal:** Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer). **Refillable Container Disposal:** For disposal, this empty container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose. **Disposal of Unused, Unwanted Product:** For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for cleanup of spills.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

FOISON Glufosinate Ammonium 200 SN

GROUP

10

HERBICIDE

FOR SALE FOR USE ON GLUFOSINATE AMMONIUM TOLERANT FIELD CORN AND ON GLUFOSINATE AMMONIUM TOLERANT SOYBEANS IN CANADA AND ON GLUFOSINATE AMMONIUM TOLERANT CANOLA IN EASTERN CANADA AND BRITISH COLUMBIA (INCLUDING THE PEACE RIVER REGIONS)

**SOLUTION
COMMERCIAL**



CAUTION - EYE IRRITANT

REGISTRATION NUMBER: 33386 PEST CONTROL PRODUCTS ACT

ACTIVE INGREDIENT: glufosinate ammonium 200 g/L

NET CONTENTS: 1L – 1,000L

READ THE LABEL AND THIS BOOKLET BEFORE USING

Product Information: +1-519-702-2185

Jiangsu Good Harvest-Weien Agrochemical Co., Ltd.

Laogang, Qidong City, Jiangsu 226221 China

Canadian Representative: Hanson Industry Company

569 Rosehill Ave., Stittsville, ON K2S 0E8

Tel: +1-519-702-2185

**FOISON Glufosinate
Ammonium 200 SN**

Table of Contents:

GENERAL INFORMATION

The Product

Section Number:

1

SAFETY AND HANDLING

Precautions, Protective Clothing
and Equipment, and
Re-entry Restriction

2

First Aid and
Toxicological Information

3

Environmental Precautions

4

Storage

5

Disposal

6

DIRECTIONS FOR USE

Crops, Weeds, Rates, Timing

7

Tank Mixes

8

Resistance Management

Recommendations

9

Application Instructions and Cautions

10

Mixing Instructions

11

Sprayer Cleanup

12

Livestock Feeding and Preharvest

Intervals

13

Notices

14

GENERAL INFORMATION

Section 1: The Product

FOISON Glufosinate Ammonium 200 SN provides control of a broad spectrum of grassy and broadleaf weeds in Canola varieties, Corn hybrids and Soybean varieties that are specially developed to be tolerant to glufosinate ammonium.

FOISON Glufosinate Ammonium 200 SN may also be applied to glufosinate ammonium tolerant inbred Corn lines grown for seed production.

For Canola hybrid seed production, FOISON Glufosinate Ammonium 200 SN may be used to remove segregating wild type plants within the female population. See Section 7 (Crops, Weeds, Rates, Timing - Canola Hybrid Seed Production).

SAFETY AND HANDLING

Section 2: Precautions, Protective Clothing and Equipment, and, Re-entry Restrictions

PRECAUTIONS:

CAUTION: EYE IRRITANT/KEEP OUT OF REACH OF CHILDREN.

May irritate eyes.

Avoid contact with eyes, skin or clothing.

Avoid breathing spray mist.

Wash thoroughly after using and before eating, drinking or smoking.

Protective Clothing and Equipment:

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. In addition, wear protective eyewear (goggles or face shield) during mixing and loading. Wear a respirator with a NIOSH-approved organic-vapour-removing cartridge with a prefilter approved for pesticides, or a NIOSH-approved canister approved for pesticides during mixing, loading and application.

Re-entry Restriction:

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. If required, certified applicators may enter treated areas within 24 hours for short-term tasks not involving hand labour if at least 4 hours has passed since application and long-sleeved shirt, long pants, chemical-resistant coveralls, chemical-resistant footwear, socks, goggles, chemical-resistant gloves, hat and a respirator with a NIOSH-approved organic-vapour-removing cartridge with a prefilter approved for pesticides OR a NIOSH-approved canister approved for pesticides is worn. Time spent in the treated area cannot exceed 1 hour in a 12-hour period during the restricted re-entry interval.

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

Section 3: First Aid and Toxicological Information

FIRST AID:

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If on skin or clothing, take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

TOXICOLOGICAL INFORMATION:

Note to Physician: If this product is ingested, endotracheal intubation and gastric lavage should be performed as soon as possible, followed by charcoal and sodium sulfate administration.

Section 4: Environmental Precautions

TOXIC to aquatic organisms and non-target terrestrial plants. Observe buffer zones specified under Directions for Use.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

Avoid spray drift onto susceptible plants and **USE ONLY FOR RECOMMENDED PURPOSES AND AT RECOMMENDED RATES.**

Section 5: Storage

CANNOT be stored below freezing.

If stored for 1 year or longer, shake well before using.

Store the tightly closed container away from feeds, seeds, fertilizers, plants and foodstuffs.

Do not use or store in or around the home.

Keep in original container during storage.

Section 6: Disposal

Recyclable Container Disposal:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

Returnable Container Disposal: Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

Refillable Container Disposal: For disposal, this empty container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

Disposal of Unused, Unwanted Product:

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for cleanup of spills.

Disposal of Unused Spray Solution:

If any spray solution remains in the tank after spraying is finished, it should be sprayed on the perimeter of the area just sprayed, away from water supplies, ditches, and irrigation canals.

DIRECTIONS FOR USE

Section 7: Crops, Weeds, Rates, Timing

Best control will be obtained when FOISON Glufosinate Ammonium 200 SN is applied at the recommended leaf stages of the weeds.

In Canola grown in Eastern Canada and British Columbia, apply FOISON Glufosinate Ammonium 200 SN from the cotyledon stage up until, but prior to, the bolting stage of Canola.

In Corn grown in Canada, apply FOISON Glufosinate Ammonium 200 SN from the 1-8 leaf stage of the Corn plant or 5-6 visible collars (the leaf is counted once the next leaf is visible in the whorl). In Soybeans grown in Canada, apply FOISON Glufosinate Ammonium 200 SN from the cotyledon to the flowering stage of the crop.

Crop	Weed	Recommended Time (Leaf Stage)								Rate
		1	2	3	4	5	6	7	8	
CANOLA, CORN, and SOYBEANS tolerant to FOISON Glufosinate Ammonium 200SN	Crabgrass	----	----	----	----	x				1.5 L/ha
	Giant Foxtail	----	----	----	----	x				
	Green Foxtail	----	----	----	----	x				
	Proso Millet	----	----	----	----	x				
	Chickweed	----	----	----	----	-----	-----	-----	x	
	Cocklebur	----	----	----	x					
	Ragweed	----	----	----	----	x				
	Redroot Pigweed	----	----	----	----	-----	-----	-----	x	
Shepherd's- purse	----	----	----	----	-----	-----	-----	x		

Crop	Weed	Recommended Time (Leaf Stage)								Rate
		1	2	3	4	5	6	7	8	
CANOLA, CORN, and SOYBEANS tolerant to FOISON Glufosinate Ammonium 200SN	Barnyard Grass	----	----	----	----	x				2 L/ha
	Bristly Foxtail	----	----	----	x					
	Fall Panicum	----	----	----	x					
	Old Witchgrass	----	----	----	x					
	Wild Oats	----	----	----	x					
	Yellow Foxtail	----	----	----	x					
	Eastern Black Nightshade	----	----	----	-----	x				
	Lady's-thumb	----	----	----	-----	-----	x			
	Lamb's-quarters	----	----	----	-----	-----	x			
	Perennial Sowthistle	----	----	----	x					
	Ragweed	----	----	----	-----	-----	-----	x		
	Stinkweed	----	----	----	-----	-----	-----	-----	x	
	Velvetleaf	----	----	----	x					
	Wild Buckwheat	----	----	----	x					
	Wild Mustard	----	----	----	x					
	Wormseed Mustard	----	----	----	-----	-----	-----	x		

Canada Thistle and Field Bindweed: For season long suppression of Canada Thistle and Field Bindweed, apply FOISON Glufosinate Ammonium 200 SN at a rate of 2 L/ha when the weeds are in the 1-6 leaf stage.

Quackgrass: For season long suppression of Quackgrass, apply FOISON Glufosinate Ammonium 200 SN at a rate of 2.5 L/ha plus ammonium sulphate at a rate of 6 L/ha (49% solution) OR 3 kg/ha (99%) when the Quackgrass is in the 1-4 leaf stage.

Voluntary Crop Weeds: FOISON Glufosinate Ammonium 200 SN will not control volunteer crop weeds that emerge from previous plantings of glufosinate ammonium tolerant crops.

ADDITIVES: For enhanced control of Velvetleaf, add ammonium sulphate to the tank at a rate of 6 L/ha (49% solution) OR 3 kg/ha (99%).

SECOND APPLICATION: A second application of FOISON Glufosinate Ammonium 200 SN can be made to fields treated initially with up to 2.5 L/ha if new weed germination or growth is present.

Apply when the new weed growth is in the correct leaf stage and up to the maximum leaf stage of the crop.

Application after the 8-leaf stage of Corn must be made with drop nozzles as a directed spray below the whorl of the Corn plant.

Application in Canola must be made prior to the bolting stage of the Canola.

Do not apply beyond the flowering stage of Soybeans.

Do not apply more than a total of 4.5 L/ha in one season.

SPLIT APPLICATION PROGRAM: For season long control of the above listed weeds a split application program with FOISON Glufosinate Ammonium 200 SN can be utilised. The first application must be a minimum of 2 L/ha and at the proper weed stage. For the second application, the rate of 1.25 L/ha may be utilized provided FOISON Glufosinate Ammonium 200 SN is applied as soon as the second flush of weeds occurs and before the maximum leaf stage of the crop.

FOR CANOLA HYBRID SEED PRODUCTION - Two applications are required to remove the segregating wild type plants. The first application should occur when the Canola is in the 2-4 leaf stage, the second application when the Canola plants are in the 4-6 leaf stage and the third application, if necessary, may be applied to the Canola up until, but prior to, the time of bolting. All applications of FOISON Glufosinate Ammonium 200 SN for hybrid seed production should be made at 2.5 L/ha.

Section 8: Tank Mixes

Consult the label of the tank mix partner for further instructions regarding directions for use, restrictions and precautions. Certain tank mix partners may have additional restrictions for leaf staging.

For control of weeds present and/or control of later flushes of annual broadleaf weeds and/or annual grasses (see Tank Mix Directions), FOISON Glufosinate Ammonium 200 SN may be tank mixed with the following herbicides when used in Eastern Canada and British Columbia:

CROP	TANK MIX PRODUCT	RATE	DIRECTIONS
CANOLA	Centurion [®] or Select [®] + Amigo Adjuvant at 0.5% v/v	63 mL/ha	For control of Volunteer Barley and improved control of Wild Oats, apply FOISON Glufosinate Ammonium 200 SN at a rate of 2.0 – 2.5 L/ha plus Centurion or Select at 63 mL/ha with adjuvant as recommended on the Centurion or Select label. Apply when the weeds are in the 1-4 leaf stage with a maximum of 2 tillers.
CORN	AAtrex Liquid 480 Herbicide*	1.75 – 2.33 L/ha	Mix with the appropriate rate of FOISON Glufosinate Ammonium 200 SN for the targeted weed species. Adjust atrazine rate to the level of residual activity desired. This tank mix will not control later flushes of triazine resistant weeds.
CORN	Banvel [®]	625 mL/ha	Mix with the appropriate rate of FOISON Glufosinate Ammonium 200 SN for the targeted weed species. This tank mix will provide enhanced control of annual broadleaf weeds.
CORN	Distinct [®]	285 g/ha	Mix with the appropriate rate of FOISON Glufosinate Ammonium 200 Herbicide for the targeted weed species. This tank mix will provide enhanced control of labeled annual broadleaf weeds.
CORN	Marksman [®]	2.5 - 3.7 L/ha	Mix with the appropriate rate of FOISON Glufosinate Ammonium 200 SN for the targeted weed species. Adjust Marksman rate to the level of residual activity desired. This tank mix will provide residual control of annual grasses and annual broadleaf weeds.
SOYBEAN	Pursuit [®]	312 mL/ha	Mix with the appropriate rate of FOISON Glufosinate Ammonium 200 SN for the targeted weed species. This tank mix will provide residual control of annual grasses and annual broadleaf weeds.

SOYBEAN	Basagran® Forte	1.75 L/ha	Mix with the appropriate rate of FOISON Glufosinate Ammonium 200 SN for the targeted weed species. This tank mix will provide enhanced control of annual broadleaf weeds.
---------	--------------------	-----------	---

*for atrazine products with different concentration, adjust the rate accordingly.

Note: Do not tank mix FOISON Glufosinate Ammonium 200 SN with fertilizers or any other chemical additives unless recommended on this label.

Section 9: Resistance Management Recommendations

For resistance management, FOISON Glufosinate Ammonium 200 SN is a Group 10 herbicide. Any weed population may contain or develop plants naturally resistant to FOISON Glufosinate Ammonium 200 SN and other Group 10 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of FOISON Glufosinate Ammonium 200 SN or other Group 10 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible, by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact +1-519-702-2185.

Section 10: Application Instructions and Cautions

Application Instructions:

- For best results, apply to emerged, young, actively growing weeds. Weeds that emerge after application will not be controlled.
- FOISON Glufosinate Ammonium 200 SN will have an effect on weeds that are larger than the recommended leaf stage, however speed of activity and control may be reduced.
- Apply FOISON Glufosinate Ammonium 200 SN in a minimum of 110 L/ha of water, at a pressure of 175-275 kPa and at a ground speed of 6-10 kph. The use of 80° or 110° flat fan nozzles is highly recommended for optimum spray coverage and canopy penetration.
- Application of the spray at a 45° angle forward will result in better spray coverage. Follow directions elsewhere on the label for the correct rate and timing of application.
- The speed of action of FOISON Glufosinate Ammonium 200 SN is influenced by environmental factors.
- At cool temperatures (below 10°C), poor moisture and low humidity, speed of action may be reduced. Generally, visual symptoms appear 2-4 days after application.
- IF RAINFALL OCCURS WITHIN 4 HOURS OF APPLICATION, EFFECTIVENESS MAY BE REDUCED.
- Weed control may be reduced when heavy dew, fog or mist are present at the time of application.
- FOISON Glufosinate Ammonium 200 SN works primarily as a contact herbicide. Thorough coverage of the weeds to be controlled is essential.
- As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.
- DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Application Cautions:

Field sprayer application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium classification. Boom height must be 60 cm or less above the crop or ground.

GROUND APPLICATION ONLY. DO NOT apply using aerial application equipment

Do not use flood jet nozzles, controlled droplet application equipment or air-assisted spray equipment. Uniform, thorough spray coverage is important to achieve consistent weed control.

DO NOT apply if winds exceed 16 kph when using open boom sprayers. DO NOT apply if winds exceed 25 kph when using hooded sprayers.

Avoid contact with desirable plants or crops either from direct application or from spray drift as severe damage may occur.

Plant back Intervals:

- 70 days for Buckwheat, Barley, Millet, Oats, Rye, Sorghum, Triticale and Wheat
- 120 days for all other crops except Field Corn, Canola and Soybeans. No plant back interval is required for Field Corn, Canola or Soybeans.

Application of FOISON Glufosinate Ammonium 200 SN to non-tolerant crops will result in severe crop injury or death of the crop.

SUSCEPTIBLE WEEDS: FOISON Glufosinate Ammonium 200 SN has an effect on all weeds and crops except for those crops which are developed to be tolerant to applications of FOISON Glufosinate Ammonium 200 SN, including when they become voluntary weeds.

Buffer zones:

Spot treatments using hand-held equipment do not require a buffer zone.

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands).

Method of Application	Crop	Buffer Zones (metres) Required for the Protection of:		
		Freshwater Habitat of Depths:		Terrestrial Habitat
		Less than 1 m	Greater than 1 m	
Field Sprayer	Corn, Canola, Soybean	1	0	1

For tank mixes, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank-mix partners.

Section 11: Mixing Instructions

FOISON Glufosinate Ammonium 200 SN must be applied with properly calibrated, clean equipment. FOISON Glufosinate Ammonium 200 SN is specially formulated to mix readily in water.

Prior to adding FOISON Glufosinate Ammonium 200 SN to the spray tank, ensure that the spray tank is thoroughly clean (see Section 12 “SPRAYER CLEANUP”).

1. Fill tank to three-quarters full with clean water prior to adding FOISON Glufosinate Ammonium 200 SN.
2. Add the correct amount of FOISON Glufosinate Ammonium 200 SN.
3. Add the remaining amount of water, begin agitation, and spray out immediately.

Note: With the exception of use with CENTURION or SELECT, when tank mixing FOISON Glufosinate Ammonium 200 SN always add the tank mix partner to the tank first, ensure that it is thoroughly mixed, THEN add FOISON Glufosinate Ammonium 200 SN. When using ammonium sulphate, ALWAYS add it to the tank first, prior to the addition of the tank mix partner and FOISON Glufosinate Ammonium 200 SN. Refer to the following specific instructions for mixing with CENTURION or SELECT.

FOISON Glufosinate Ammonium 200 SN and CENTURION or SELECT: Glufosinate Ammonium Tolerant Canola only

When tank mixing FOISON Glufosinate Ammonium 200 SN and CENTURION or SELECT, always add AMIGO adjuvant to the tank first, *then* add FOISON Glufosinate Ammonium 200 SN to the tank followed by the CENTURION or SELECT.

The addition of an anti-foaming agent may reduce foaming, especially when using soft water.

1. Thoroughly clean the sprayer by flushing the system with water containing detergent (see Section 12, “Sprayer Cleanup”).
2. Fill clean spray tank half full with clean water. Start agitation system.
3. Add the correct amount of AMIGO. Continue to agitate until AMIGO is thoroughly mixed.
4. STOP agitation. Add the required amount of FOISON Glufosinate Ammonium 200 SN and Crop Desiccant MP to the spray tank. Start agitation system.
5. Add the correct amount of CENTURION or SELECT along with the remaining amount of water necessary to fill the spray tank.
6. Continue to agitate or run the by-pass system and spray out immediately.
7. After any break in the spraying operation, agitate thoroughly before spraying again. Check inside the tank to ensure that sprayer agitation is sufficient to remix the spray materials. Do not allow the mixture to sit overnight.
8. If an oil film starts to build up in the tank, drain it and clean tank with strong detergent solution.
9. Immediately after use, thoroughly clean the sprayer by flushing the system with water containing detergent (see Section 12, “Sprayer Cleanup”).

Note: The addition of an anti-foaming agent may reduce foaming, especially when using soft water.

Note: Ensure that all circuits (pipes, booms, etc.) have the correct FOISON Glufosinate Ammonium 200 SN/water concentration before application is started.

Section 12: Sprayer Cleanup

Before and after using FOISON Glufosinate Ammonium 200 SN always complete a thorough cleaning of the spray tank, lines and filter. Spray equipment should be thoroughly rinsed using a strong detergent solution.

Section 13: Livestock Feeding and Preharvest Intervals (PHIs)

CANOLA:

Grain and meal from treated Canola can be fed to livestock.

DO NOT graze or feed other portions of the treated Canola to livestock; there are not sufficient data to support such use.

When FOISON Glufosinate Ammonium 200 SN is tank mixed with CENTURION or SELECT, observe a PHI of 60 days from the date of treatment (or last treatment when a second application has been made).

CORN:

DO NOT apply within 86 days of harvesting Corn grain.

DO NOT graze the treated fields within 20 days of application.

SOYBEAN:

DO NOT apply within 70 days of harvesting Soybean seed.

DO NOT graze the treated fields within 20 days of application.

Section 14: Notices

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.